


O(1) GeV DM in SUSY and a very light pseudoscalar at the LHC

Doyoun Kim
Asia Pacific Center for Theoretical Physics

in collaboration with
C. Han, S. Munir and M. Park
arXiv:1504.05085
JHEP 1507 (2015) 002

Higgs is found. Then what else?



- 125 GeV is quite a safe mass if we consider SM as the effective theory below 100 ~ 1000 TeV.

Higgs is found. Then what else?

- 125 GeV is quite a safe mass if we consider SM as the effective theory below 100 ~ 1000 TeV.
- And LHC has so far seen no remarkable observation to be regarded as a new physics signal.
- But still we keep our positions and waiting for some news.

Higgs is found. Then what else?

- Attempts to survey the structure of our vacuum or universe are going on.
- Hierarchy Problem - SUSY, Composite Higgs, Twin Higgs, Extra D, etc.
- Dark Matter
- Neutrino Oscillation (Nobel Prize 2015)
- Other Precision measurements: B-physics, neutron EDM, muon $g-2$, etc.

Higgs is found. Then what else?

- Attempts to survey the structure of our vacuum or universe are going on.

- Hierarchy Problem - SUSY, Composite Higgs, Twin Higgs, Extra D, etc.

- Dark Matter

- Neutrino Oscillation

- Other Precision measurements: B-physics, neutron EDM, muon $g-2$, etc.

SUSY Model of SM particles

- Theoretically sound and predictive model
- Rich in phenomenological implications:
3 neutral Higgs, DM candidate (LSP neutralino with R-parity)
- However, no evidence at all in the LHC Run I.
- Because m_h^2 requires a large radiative correction from the stops, and it implies $m_{\text{soft}} \sim \text{TeV}$ or X_t (A_t) $\sim \text{multi TeV}$.

SUSY Model of SM particles

- Theoretically sound and predictive model
- Rich in phenomenological implications:
3 neutral Higgs, DM candidate (LSP neutralino with R-parity)
- However, no evidence at all in the LHC Run I.
- Because m_h^2 requires a large radiative correction from the stops, and it implies $m_{\text{soft}} \sim \text{TeV}$ or X_t (A_t) $\sim \text{multi TeV}$.
- Large fine-tuning between $m_{H_i}^2, \mu$ and M_Z .

$$\frac{M_Z^2}{2} = \frac{m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2$$

SUSY Model of SM particles


- Theoretically sound and predictive model
- Rich in phenomenological implications:
3 neutral Higgs, DM candidate (LSP neutralino with R-parity)
- However, no evidence at all in the LHC Run I.
- Because m_h^2 requires a large radiative correction from the stops, and it implies $m_{\text{soft}} \sim \text{TeV}$ or X_t (A_t) $\sim \text{multi TeV}$.
- Large fine-tuning between $m_{H_i}^2, \mu$ and M_Z .
- What are the issues in the Dark Matter side?

DM might be just around the corner of Higgs

- SUSY suggests a good DM candidate (LSP with R-Parity).


DM might be just around the corner of Higgs

- MSSM neutralino covers a large range of DM mass.



DM might be just around the corner of Higgs

- SUSY suggests a good DM candidate (LSP with R-Parity).




DM might be just around the corner of Higgs

- SUSY suggests a good DM candidate (LSP with R-Parity).
- If we consider the relic abundance measured in PLANCK exp., an MSSM wino-higgsino mixture is promising in a wide mass range.
- Bino DM is also possible when it annihilates resonantly via some mediators such as Z' , Higgs, etc.
- Or we can suggest a more complicated model to prevent our universe from over-closure. (Bino is an NLSP and we have lighter one such as axino DM.)
- From MSSM, it is hard to accommodate the neutralino DM below 10 GeV mass range.


DM might be just around the corner of Higgs

- SUSY suggests a good DM candidate (LSP with R-Parity).
- But the DM search is getting challenged more.


DM might be just around the corner of Higgs



DM might be just around the corner of Higgs




DM might be just around the corner of Higgs



How to realize light DM in SUSY

- It is difficult to realize an O(1) GeV DM in the MSSM
-> But in the NMSSM, it is possible. (1504.05085)



NMSSM setup for singlino DM

- Next-to-Minimal Supersymmetric extension of the Standard Model
- Introduces a SM singlet superfield to reduce the fine-tuning of the Higgs mass. And all mass scales are introduced at the SUSY breaking scale.

$$\begin{array}{ccc}
 W \supset \mu H_u H_d & \xrightarrow[\text{NMSSM}]{\text{MSSM}} & W \supset \lambda S H_u H_d + \frac{\kappa}{3} S^3 \\
 V \supset B \mu h_u h_d & & V \supset \lambda A_\lambda S h_u h_d
 \end{array}$$

- SM Higgs mass is lifted up at tree level by a sizable interaction with singlet sector.

$$m_{H_{1,2}}^2 \simeq \frac{1}{2} \left\{ m_Z^2 + 4(\kappa s)^2 - \kappa s A_\kappa \right. \\
 \left. \mp \sqrt{[m_Z^2 - 4(\kappa s)^2 - \kappa s A_\kappa]^2 + 4\lambda^2 v^2 [2\lambda s - (A_\lambda + \kappa s) \sin 2\beta]^2} \right\}$$

NMSSM setup for singlino DM

- Next-to-Minimal Supersymmetric extension of the Standard Model
- Introduces a SM singlet superfield to reduce the fine-tuning of the Higgs mass. And all mass scales are introduced at the SUSY breaking scale.

$$\begin{array}{ccc}
 W \supset \mu H_u H_d & \xrightarrow[\text{NMSSM}]{\text{MSSM}} & W \supset \lambda S H_u H_d + \frac{\kappa}{3} S^3 \\
 V \supset B \mu h_u h_d & & V \supset \lambda A_\lambda S h_u h_d
 \end{array}$$

- Light singlino-like DM is decoupled from the higgsino states. ($\kappa s \ll \mu_{\text{eff}}$)

$$\begin{pmatrix}
 m_{\tilde{B}, \tilde{W}} & m_{\tilde{H}_{u,d}} & m_{\tilde{S}} \\
 \dots & & \\
 & 0 & -\mu_{\text{eff}} & -\lambda v_u \\
 & -\mu_{\text{eff}} & 0 & -\lambda v_d \\
 & -\lambda v_u & -\lambda v_d & 2\kappa s
 \end{pmatrix}$$

NMSSM setup for singlino DM

- Next-to-Minimal Supersymmetric extension of the Standard Model
- Introduces a SM singlet superfield to reduce the fine-tuning of the Higgs mass. And all mass scales are introduced at the SUSY breaking scale.

$$\begin{array}{ccc}
 W \supset \mu H_u H_d & \xrightarrow[\text{NMSSM}]{\text{MSSM}} & W \supset \lambda S H_u H_d + \frac{\kappa}{3} S^3 \\
 V \supset B \mu h_u h_d & & V \supset \lambda A_\lambda S h_u h_d
 \end{array}$$

- Light singlino-like DM is decoupled from the higgsino states. ($\kappa s \ll \mu_{\text{eff}}$)
- Strong mass correlation between $m_{\tilde{\chi}_1^0}$ and $m_{A_1^0}$ ($\sim \kappa s$) for a large $\tan \beta$.

$$m_{A_1^0}^2 \simeq \lambda(A_\lambda + 4\kappa s) \frac{v^2 \sin^2 2\beta}{2s} - 3\kappa s A_\kappa$$

NMSSM setup for singlino DM

- Next-to-Minimal Supersymmetric extension of the Standard Model
- Introduces a SM singlet superfield to reduce the fine-tuning of the Higgs mass. And all mass scales are introduced at the SUSY breaking scale.

$$\begin{array}{ccc} W \supset \mu H_u H_d & \xrightarrow[\text{NMSSM}]{\text{MSSM}} & W \supset \lambda S H_u H_d + \frac{\kappa}{3} S^3 \\ V \supset B \mu h_u h_d & & V \supset \lambda A_\lambda S h_u h_d \end{array}$$

- Light singlino-like DM is decoupled from the higgsino states. ($\kappa S \ll \mu_{\text{eff}}$)
- Strong mass correlation between $m_{\tilde{\chi}_1^0}$ and $m_{A_1^0}$ ($\sim \kappa S$) for a large $\tan \beta$.
 \Rightarrow Light singlino and light pseudo-scalar can be easily realized.

Parameter Scan and Benchmark Points

- Relic abundance consistent with PLANCK 2015 data.
 - ⇒ Necessity to enhance the annihilation channel of LSP
 - ⇒ Resonant s-channel process: $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow A_1^0 \rightarrow f \bar{f}$
 H_1^0 : p-wave suppressed
 $2m_{\tilde{\chi}_1^0} = m_{A_1^0} \sim 3 \text{ GeV}$
- Rare B meson decay constraints.
- LEP Electroweak Precision Test.
- Etc.


Parameter Scan and Benchmark Points

- Parameter sets we have found interesting

	BP1	BP2
<i>Model parameters</i>		
M_0 (GeV)	1951.1	1826.0
$M_{1/2}$ (GeV)	892.24	929.2
A_0 (GeV)	2462.2	2626.2
μ_{eff} (GeV)	191.34	164.52
$\tan \beta$	14.056	19.785
λ	0.0814	0.3102
κ	0.0002	0.0008
A_λ (GeV)	4080.2	3596.7
A_κ (GeV)	-3.6681	-6.8466

LHC search


- How to optimally survey BPs in the collider:
BP1 and 2 have distinguishable features at the LHC (300/fb).



	BP1	BP2
<i>Masses</i>		
$m_{\tilde{\chi}_1^0}$ (GeV)	1.0025	1.4081
$m_{\tilde{\chi}_2^0}$ (GeV)	189.09	170.13
$m_{\tilde{\chi}_3^0}$ (GeV)	-201.67	-182.27
$m_{\tilde{\chi}_1^\pm}$ (GeV)	194.97	167.72
m_{A_1} (GeV)	2.1776	2.9856
m_{H_2} (GeV)	124.12	125.79
<i>Branching Ratios</i>		
$BR(\tilde{\chi}_2^0 \rightarrow Z \tilde{\chi}_1^0)$	0.634	0.603
$BR(\tilde{\chi}_2^0 \rightarrow A_1 \tilde{\chi}_1^0)$	0.004	0.089
$BR(\tilde{\chi}_3^0 \rightarrow Z \tilde{\chi}_1^0)$	0.736	0.704
$BR(\tilde{\chi}_3^0 \rightarrow A_1 \tilde{\chi}_1^0)$	0.004	0.081
$BR(A_1 \rightarrow \mu^+ \mu^-)$	0.039	0.087

LHC search


- How to optimally survey BPs in the collider
BP1 and 2 have distinguishable features at the LHC (300/fb).




	BP1	BP2
<i>Masses</i>		
$m_{\tilde{\chi}_1^0}$ (GeV)	1.0025	1.4081
$m_{\tilde{\chi}_2^0}$ (GeV)	189.09	170.13
$m_{\tilde{\chi}_3^0}$ (GeV)	-201.67	-182.27
$m_{\tilde{\chi}_1^\pm}$ (GeV)	194.97	167.72
m_{A_1} (GeV)	2.1776	2.9856
m_{H_2} (GeV)	124.12	125.79
<i>Branching Ratios</i>		
$BR(\tilde{\chi}_2^0 \rightarrow Z\tilde{\chi}_1^0)$	0.634	0.603
$BR(\tilde{\chi}_2^0 \rightarrow A_1\tilde{\chi}_1^0)$	0.004	0.089
$BR(\tilde{\chi}_3^0 \rightarrow Z\tilde{\chi}_1^0)$	0.736	0.704
$BR(\tilde{\chi}_3^0 \rightarrow A_1\tilde{\chi}_1^0)$	0.004	0.081
$BR(A_1 \rightarrow \mu^+\mu^-)$	0.039	0.087

LHC search

- How to optimally survey BPs in the collider:
BP1 and 2 have distinguishable features at the LHC (300/fb).



2.7 σ significance level expected at the LHC Run II



S/B ~ 15 expected at the LHC Run II

Summary

- Higgs mass implies that the simplest implementation of SUSY still requires a fine-tuning. NMSSM may help to ameliorate the tension.
- DM search calls for back-up by the LHC data.
- NMSSM accommodates $O(1)$ GeV DM via the decoupled singlet sector, and sometimes produces the collimated objects which is easily missed out.
- A proper treatment for them may catch priceless events at the LHC Run II.
- Especially, a collimated soft final states are useful for the model with a compressed mass spectrum. [1502.03734]
- Thank you.

BACKUP SLIDES

LHC search

- Similar to and Motivated by the lepton-jet object.
- Define a collimated muon system as a single detector object.
 1. Require the transverse momentum > 10 GeV for each muon.
Invariant mass of the muon pair < 5 GeV
 2. Scalar sum of transverse momenta of all additional charged tracks inside the jet cone size $\Delta R = 0.4$, less than 3 GeV.
- Main Backgrounds, containing two collinear muons along with a third lepton and MET, include $W(\rightarrow l^\pm \nu)\gamma^*$ and $Z(\rightarrow l^+ l^-)\gamma^*$.

LHC search

- Final Result

	BP1	$W\gamma^*$	$Z\gamma^*$	$Wb\bar{b}$
Cross section (fb)	0.178	246.9	10.0	3770.0
Cut efficiency	0.123	2.15×10^{-4}	6×10^{-5}	1×10^{-6}
Effective cross section (fb)	0.022	0.053	0.0006	0.003
No. of events	6.6	15.9	0.18	0.9

	BP2	$W\gamma^*$	$Z\gamma^*$	$Wb\bar{b}$
Cross section (fb)	3.93	246.9	10.0	3770.0
Cut efficiency	0.050	5.3×10^{-5}	3×10^{-5}	1×10^{-6}
Effective cross section (fb)	0.197	0.013	0.0003	0.003
No. of events	59.1	3.9	0.09	0.9

Point	S/B in analysis		\mathcal{Z} (σ) in analysis	
	3ℓ (SRZc region)	μ_{col}	3ℓ (SRZc region)	μ_{col}
BP1	0.591	0.42	2.7	1.2
BP2	0.436	15	2.0	27